


State Math Contest – Junior Exam

SOLUTIONS

8. The large triangle below is divided into two triangles of areas α and β . Find α/β .



- a) $\frac{3}{5}$ b) $\frac{1}{2}$ c) $\frac{4}{6}$
d) $\frac{3}{4}$ e) $\frac{7}{11}$

Solution:

Correct answer: $\frac{3}{5}$ (a)
Over bases 3 and 5, the two triangles have the same altitude.

9. Let

$$f(x) = \begin{cases} 2x & \text{for } 0 \leq x \leq 0.5 \\ 2(1-x) & \text{for } 0.5 < x \leq 1. \end{cases}$$

If $x_0 = \frac{6}{7}$ and $x_n = f(x_{n-1})$ for $n \geq 1$, find x_{100} .

- a) $\frac{2}{7}$ b) $\frac{4}{7}$ c) $\frac{6}{7}$
d) $\frac{10}{7}$ e) $\frac{96}{7}$

Solution:

Correct answer: $\frac{2}{7}$ (a)
 $x_0 = 6/7, x_1 = 2/7, x_2 = 4/7, x_3 = 6/7, x_4 = 2/7, x_5 = 4/7$.
The sequence repeats every three terms.
 $x_{100} = x_{97} = \dots = x_1 = 2/7$.

10. When $3x^{12} - x^3 + 5$ is divided by $x + 1$ the remainder is:

- a) 1 b) 3 c) 5
d) 7 e) 9

Solution:

Correct answer: 9 (e)
By the Remainder Theorem, the answer is the polynomial evaluated at $x = -1$.
 $3(-1)^{12} - (-1)^3 + 5 = 3 + 1 + 5 = 9$.

13. High schools in Utah are divided into different classifications, with the largest schools classified as 5A and the next largest schools classified as 4A.

If the smallest 5A school is reclassified as 4A, what will happen to the average size of schools in the two classifications?

- a) The average 4A size will go down, and the average 5A size will go up.
- b) The average 4A size will go up, and the average 5A size will go up.
- c) The average 4A size will go down, and the average 5A size will go down.
- d) The average 4A size will go up, and the average 5A size will go down.
- e) None of the above is always true.

Solution:

Both averages will go up. (b)

14. How many ways can you write 5 as the sum of one or more positive integers if different orders are not counted differently? For example, there are three ways to write 3 in this way: $1 + 1 + 1$, $1 + 2$, and 3.

- a) 7
- b) 6
- c) 8
- d) 5
- e) 10

Solution:

Correct answer: 7 (a)
5,
 $4 + 1$,
 $3 + 2$,
 $3 + 1 + 1$,
 $2 + 2 + 1$,
 $2 + 1 + 1 + 1$,
 $1 + 1 + 1 + 1 + 1$.

15. How many real solutions does the equation $x^{3/2} - 32x^{1/2} = 0$ have?

- a) 0
- b) 1
- c) 2
- d) 3
- e) 4

Solution:

Correct answer: 2 (c)
Let $y = x^{1/2} = \sqrt{x}$. the original equation becomes $y^3 - 32y = 0$ or $y(y^2 - 32) = 0$. Thus $y = 0$ or $y = \pm\sqrt{32} = \pm 4\sqrt{2}$. If $y = \sqrt{x} = 0$, then $x = 0$. If $y = \sqrt{x} = \sqrt{32}$, then $x = 32$. There is no real value of x for which $y = \sqrt{x} = -\sqrt{32}$, since for x real $\sqrt{x} \geq 0$. So there are two real solutions to the equation.

25. If $|r| < 1$, then $(a)^2 + (ar)^2 + (ar^2)^2 + (ar^3)^2 + \dots =$

a) $\frac{a^2}{(1-r)^2}$

b) $\frac{a^2}{1+r^2}$

c) $\frac{a^2}{1-r^2}$

d) $\frac{4a^2}{1+r^2}$

e) none of these

Solution:

Correct answer: $\frac{a^2}{1-r^2}$ (c)

This is a geometric series with first term a^2 and ratio r^2 . The sum of an infinite geometric series with ratio of absolute value less than 1 is the first term divided by one minus the ratio; i.e., $\frac{a^2}{1-r^2}$.

26. Consider the set of colors {white, black, red, orange, yellow, green, blue, purple}. We define the operation of addition (+) on this set of colors such that if two colors from the set are added together, we obtain another (not necessarily distinct) color in the set. For example, the following rules are always satisfied.

$$\begin{aligned} \text{blue} + \text{red} &= \text{purple} \\ \text{blue} + \text{yellow} &= \text{green} \\ \text{yellow} + \text{red} &= \text{orange} \\ \text{red} + \text{blue} + \text{yellow} &= \text{white} \end{aligned}$$

In addition if black is added to another color, we obtain that color, while if a color is added to itself we obtain black. If this addition is commutative and associative, fill in the blank such that

$$\text{yellow} + \text{green} + \underline{\hspace{2cm}} = \text{purple}$$

is a true statement.

a) Green

b) Red

c) Black

d) Blue

e) Purple

Solution:

Correct answer: Red (b)

yellow + green = yellow + yellow + blue
= black + blue = blue

We know red + blue = purple.

27. When buying a bike from the *Math Bikes* company, there are three extra options to choose (a bell, a rear fender, and a basket), each of which you can choose to add to the bike or choose not to add it. If *Math Bikes* has sold 300 bikes, what is the largest number of bikes that you can guarantee to have exactly the same extras as each other?

a) 8

b) 37

c) 38

d) 43

e) 292

Solution:

Correct answer: 38 (c)

There are $2^3 = 8$ possible bike types. Now $300 \div 8 = 37.5$. If there were 37 or less of each type of bike, then there would be less than 300 bikes. So there must be at least 38 bikes that have exactly the same extras.

28. A square and an equilateral triangle have the same area. Let A be the area of the circle circumscribed around the square and B be the area of the circle circumscribed around the triangle. Find $\frac{A}{B}$.

a) $\frac{3\sqrt{3}}{8}$

b) $\frac{3\sqrt{3}}{6}$

c) $\frac{3\sqrt{3}}{4}$

d) $\frac{3}{8}$

e) $\frac{3}{4}$

Solution:

Correct answer: $\frac{3\sqrt{3}}{8}$ (a)

The \mathcal{A} be the common area. Let e be the edge length of the square, then $\mathcal{A} = e^2$. Let s be the side length of the triangle, then $\mathcal{A} = \frac{s^2\sqrt{3}}{4}$.

The center of the square is the point where the two diagonals meet. The radius r of the circle circumscribed around the square is the distance from the center of the square to the four vertices which is $\frac{e}{\sqrt{2}}$.

The circumcenter of the triangle is the point where the perpendicular bisectors meet which, in this case, is the same as the centroid or the point where the medians meet. The centroid lies on the medians two-thirds the distance from the vertex to the midpoint of the opposite side. The length of a median is $\frac{s\sqrt{3}}{2}$. So the radius R of the circle circumscribed

about the triangle is $\frac{2}{3} \cdot \frac{s\sqrt{3}}{2} = \frac{s}{\sqrt{3}}$

$$\frac{A}{B} = \frac{\pi r^2}{\pi R^2} = \frac{r^2}{R^2} = \frac{e^2}{2} \div \frac{s^2}{3} = \frac{\mathcal{A}}{2} \div \frac{4\mathcal{A}}{3\sqrt{3}} = \frac{3\sqrt{3}}{8}.$$

31. For a certain baseball team the probability of winning any game is P , (the probability of winning a particular game is independent of any other games). What is the probability the team wins 3 out of 5 games?

- a) $10P^2(1 - P)^3$ b) $10P^3(1 - P)^2$ c) $5P^3(1 - P)^2$
d) $5P^2(1 - P)^3$ e) $P^3(1 - P)^2$

Solution:

Correct answer: $10P^3(1 - P)^2$ (b)

The number of possible outcomes from playing 5 games is 2^5 . There are 5 choose 3 or 10 ways of winning 3 games and losing 2; e.g. winning the first three and losing the last two, winning the first two and fourth game and losing the other two, etc. Each of these ways of winning 3 and losing 2 games has probability $P^3(1 - P)^2$. So the total probability of winning 3 games and losing 2 is $10P^3(1 - P)^2$.

32. If x is the *fraction* of numbers between 1 and 1,000, inclusive, which contain 4 as a digit, and y is the *fraction* of numbers between 1 and 10,000, inclusive which contain 4 as a digit, what is x/y ?

- a) $2/3$ b) $3/4$ c) $27/34$
d) $271/3439$ e) $2710/3439$

Solution:

Correct answer: $2710/3439$ (e)

We can count how many numbers in $\{1, 2, \dots, 1,000\}$ contain 4 as a digit using the inclusion exclusion principle: If one digit is a 4, there are 10^2 possibilities for the other three digits, and we can do this allowing the 4 to be in the ones, the tens, and the hundreds digit, but we've overcounted. We need to subtract off the number of ways to get *two* 4's. But then we've undercounted, and we need to add back the number of ways to get *three* 4's:

$$3 \cdot (100) - 3 \cdot (10) + 1 = 271.$$


So, $x = 271/1,000$. We calculate y similarly: there are

$$4 \cdot (1,000) - 6(100) + 4(10) - 1 = 3439$$

numbers in $\{1, 2, \dots, 10,000\}$ containing 4 as a digit, so $y = 3439/10,000$. Thus,

$$\frac{x}{y} = \frac{\frac{271}{1,000}}{\frac{3439}{10,000}} = \frac{2710}{3439}.$$

33. Given that the area of the outer circle is ten square units, find the area of any one of the three equal circles which are tangent to each other and to the outer circle, and inscribed inside the circle of ten square units.



- a) $30(7 - 4\sqrt{3})$ square units. b) 2.5 square units.
c) $\frac{10}{(3 + \sqrt{2})}$ square units. d) $\sqrt[3]{10}$ square units.
e) 2 square units.

Solution:

Correct answer: $30(7 - 4\sqrt{3})$ square units (a)
The radius of the outer circle is $R = r + r \csc(\pi/3)$ where r is the radius of an inner circle.

34. Let $f(x) = 9x^2 + dx + 4$. For certain values of d , the equation $f(x) = 0$ has only one solution. For such a value of d , which value of x could be a solution to $f(x) = 0$?

- a) $\frac{2}{3}$ b) 1 c) $\frac{4}{3}$
d) 3 e) 12

Solution:

Correct answer: $\frac{2}{3}$ (a)
Using the quadratic equation gives $d = \pm 12$ as the only values for one real solution.

35. The natives of Wee-jee Islands rate 2 spears as worth 3 fishhooks and a knife, and will give 25 coconuts for 3 spears, 2 knives, and a fishhook together. Assuming each item is worth a whole number of coconuts, how many coconuts will the natives give for each article separately?

	Item	Worth in Coconuts
a)	fishhook	1
	knife	3
	spear	3

	Item	Worth in Coconuts
b)	fishhook	2
	knife	2
	spear	4

	Item	Worth in Coconuts
c)	fishhook	1
	knife	5
	spear	4

	Item	Worth in Coconuts
d)	fishhook	3
	knife	3
	spear	6

	Item	Worth in Coconuts
e)	fishhook	2
	knife	4
	spear	5

Solution:

Correct Ans: (e)

Let S , F , and K be the value of a spear, knife and fishhook in coconuts. Then

$$2S = 3F + K$$

and

$$25 = 3S + 2K + F.$$

Multiply the second equation above by 2 and the first equation by 3 to find

$$50 = 6S + 4K + 2F$$

$$6S = 9F + 3K.$$

Substituting the second equation into the first gives:

$$50 = 11F + 7K.$$

We know that F and K are nonnegative integers. If $F \geq 5$ then $11F + 7K \geq 55$, hence $F = 0, 1, 2, 3$, or 4 . We try each of these four cases, and only $F = 2$, $K = 4$ works, and in this case $S = 5$.

36. An octagon in the plane is symmetric about the x -axis, the y -axis, and the line whose equations is $y = x$. If $(1, \sqrt{3})$ is a vertex of the octagon, find its area.

a) $6\sqrt{3}$

b) 11

c) $6 + 2\sqrt{3}$


d) $2 + 6\sqrt{3}$

e) $4 + 4\sqrt{3}$

Solution:

Correct answer: $4 + 4\sqrt{3}$ (e)

The vertices of the octagon are $(\pm 1, \pm\sqrt{3})$ and $(\pm\sqrt{3}, \pm 1)$. The area can be computed by computing the area of the enclosing square that is $2\sqrt{3} \times 2\sqrt{3}$ and subtracting the area of the four right isosceles triangles or, equivalently, the area of two squares that are $\sqrt{3} - 1 \times \sqrt{3} - 1$. The area is $(2\sqrt{3})^2 - 2(\sqrt{3} - 1)^2 = 12 - 2(4 - 2\sqrt{3}) = 4 + 4\sqrt{3}$.



37. Five points are placed in a square with side length 1. What is the largest distance d so that every pair of points is at least d apart from each other?

a) 1

b) $\sqrt{2}$

c) $\sqrt{3}/2$

d) $1/2$

e) $\sqrt{2}/2$

Solution:

Answer: $\sqrt{2}/2$ (e)

If we divide the square into 4 equal squares each of side length $1/2$. Now place the 5 points among these 4 squares. One square must have two points. The farthest distance of these two points could be is $\sqrt{2}/2$.

38. A square number is an integer number which is the square of another integer.

Positive square numbers satisfy the following properties:

- The units digit of a square number can only be 0, 1, 4, 5, 6, or 9.
- The digital root of a square number can only be 1, 4, 7, or 9.

The *digital root* is found by adding the digits of the number. If you get more than one digit you add the digits of the new number. Continue this until you get to a single digit. This digit is the digital root.

One of the following numbers is a square. Which one is it?

- a) 4, 751, 006, 864, 295, 101
- b) 3, 669, 517, 136, 205, 224
- c) 2, 512, 339, 789, 576, 516
- d) 1, 898, 732, 825, 398, 318
- e) 5, 901, 643, 220, 186, 107

Solution:

Correct Ans: 2, 512, 339, 789, 576, 516 (c)
(d), and (e) are ruled out because of the first property.
(a) and (b) are ruled out because their digit roots are 5 and 8.
(c) has a digit root of 7 thus it is the square number.

39. A regular octahedron is formed by setting its vertices at the centers of the faces of the cube. Another regular octahedron is formed around the cube by making the center of each triangle of the octahedron hit at a vertex of the cube. What is the ratio of the volume of the larger octahedron to that of the smaller octahedron?

- a) $2\sqrt{2}$
- b) $27/8$
- c) $3\sqrt{3}$
- d) 8
- e) 27

Solution:

Correct answer: 27 (e)
If the cube is centered at the origin with x,y,z three dimensional coordinates, then the vertices of the cube can be at $(\pm a, \pm a, \pm a)$. The vertices of the smaller octahedron are $(\pm a, 0, 0), (0, \pm a, 0), (0, 0, \pm a)$. The vertices of the larger octahedron are $(\pm 3a, 0, 0), (0, \pm 3a, 0), (0, 0, \pm 3a)$ since the centroid of the triangle with vertices $(3a, 0, 0), (0, 3a, 0), (0, 0, 3a)$ is (a, a, a) . Since the larger octahedron is similar to the smaller by a scale factor of 3, the volume is $3^3 = 27$ times as large.

40. In $\triangle ABC$, $AC = 13$, $BC = 15$ and the area of $\triangle ABC = 84$. If $CD = 7$, $CE = 13$, and the area of $\triangle CDE$ can be represented as $\frac{p}{q}$ where p and q are relatively prime positive integers, find q .


a) 3

b) 5

c) 7

d) 11

e) 13



Solution:

Correct answer: 5 (b)

The area of $\triangle ABC$ is $\frac{1}{2} \cdot AC \cdot BC \cdot \sin C = \frac{1}{2} \cdot 13 \cdot 15 \cdot \sin C = 84$.

The area of $\triangle CDE$ is $\frac{1}{2} \cdot CD \cdot CE \cdot \sin C = \frac{1}{2} \cdot 7 \cdot 13 \cdot \sin C$. Let $[ABC]$ be the area of $\triangle ABC$ and $[CDE]$ be the area of $\triangle CDE$. Then $\frac{[CDE]}{[ABC]} = \frac{7 \cdot 13}{13 \cdot 15} = \frac{7}{15}$. So $[CDE] = \frac{7}{15}[ABC] = \frac{7}{15}84 = \frac{7 \cdot 28}{5} = \frac{196}{5}$.